
Target: Stroke[®] Phase III: Door-to-Needle within 45 Minutes



25-49%

50-75%

Iowa 151/327 (30) Massachusetts 296/661 (61) Mississippi 209/425 (46) New Mexico 72/151 (14)

Alabama 315/556 (28) Arizona 351/499 (30) Arkansas 353/686 (70) Connecticut 245/388 (24) Illinois 810/1326 (100) Indiana 451/724 (45) Kentucky 414/605 (49) Louisiana 482/650 (31) Maine 75/125 (7)

Maryland 432/726 (40) Michigan 818/1163 (75) Minnesota 436/615 (31) Missouri 889/1221 (62) Montana 80/125 (18) Nebraska 119/225 (32) Nevada 259/395 (18) New Jersey 725/1019 (55) New York 1438/2055 (129)

©2025 American Heart Association, Inc. All rights reserved Data has not been risk-adjusted. Any use without the express written consent of the American Heart Association is strictly prohibited.

76-84%

North Carolina 1080/1545 (77) North Dakota 84/147 (22) Ohio 996/1530 (87) Oklahoma 309/490 (24) Oregon 408/594 (32) Pennsylvania 1018/1726 (120) South Carolina 652/913 (49) Tennessee 981/1312 (52) Texas 2683/3597 (168)

Washington 536/808 (55) West Virginia 125/229 (17) Wisconsin 404/698 (74)



Colorado 562/717 (33) Delaware 171/218 (6) Florida 3390/4081 (175) Georgia 1138/1476 (84) Hawaii 214/270 (13) Kansas 210/264 (19) New Hampshire 115/149 (12)

Virginia 762/984 (44)

Utah 264/303 (17)